

BLOCKING INTRAMEDULLARY OSTEOSYNTHESIS - AS EFFECTIVE METHOD FOR FEMORAL FRACTURES

Hamroev Behzod Uktamovich, Akhmedov Shamshod Shavkatovich

Bukhara state medical institute, Bukhara, Uzbekistan

Abstract. The analysis of complications after surgery with use of various fixation devices testifies to the need of conducting studies aimed at the definition of indications to the choice of fixation device for treatment of distal femur fractures. In the paper the comparative data of surgical treatment results of patients with distal femur fractures by means of bone fixators with angle stability (Gr. I; n=14) and locking intramedullary rod (Gr. II; n=13) are given. On the basis of results analysis indications for the choice of fixation device are formulated and grounded.

Keywords: distal femur fracture, LCP-DF (Locking Compression Plate Distal Femur), locking retrograde intramedullary nailing.

Introduction. The treatment of periarticular and intraarticular fractures today remains one of the topical and complex issues of traumatology. The incidence of fractures of the distal femur is approximately 6–8% among all skeletal fractures and about 12–25% among fractures of the femur [1, 2, 3, 4, 5, 6, 7, 8, 10, 13, 30, 32, 34, 36, 54, 57, 59]. Despite some success in the treatment of injuries of the musculoskeletal system, from 5 to 54% of cases of femoral fractures lead to various unsatisfactory outcomes - delayed consolidation, fracture nonunion, pseudarthrosis, limb deformity, persistent knee joint dysfunction [9, 11, 12, 15, 17, 20, 25, 26, 35, 41, 43, 44, 46, 51, 52].

Today, among traumatologists, supporters of the use of a plate and an intramedullary locking rod for the treatment of fractures of the distal femur can be conditionally distinguished [14, 16, 18, 19, 21, 22, 24, 28, 53, 55, 56, 58]. At the same time, there is no clear separation of indications for the use of one or another fixator. An analysis of complications after osteosynthesis with various fixators indicates the need for studies aimed at determining the indications for choosing a

fixator in the treatment of fractures of the distal femur [23, 27, 29, 31, 33, 37, 38, 40, 42, 45, 47, 48, 49, 50].

The aim of this work is to evaluate the results of surgical treatment of patients with fractures of the distal femur, to determine the indications for choosing a fixator.

Materials and methods

A study was made of 27 patients with fresh fractures of the distal femur, who underwent surgical fixation of the fracture by submerged metal osteosynthesis in the period from 2018 to 2020.

We divided the patients into two groups depending on the type of fixator used - extraosseous or intraosseous.

- Group I included 14 patients with intra- and periarticular fractures of the femur, who underwent closed or open reposition of fracture fragments, extraosseous osteosynthesis with a plate with angular stability. In 9 cases, the MIPO technique (Minimally Invasive Plate Osteosynthesis - minimally invasive plate osteosynthesis) was used; in 5 cases, open reduction of fragments was required.

- Group II included 13 patients with intra- and periarticular fractures of the femur, who underwent closed or open reposition of fracture fragments and retrograde fixation with an intramedullary nail. In 9 cases, the technique of closed indirect reduction of fragments was used, fixation of the fracture without exposing the fracture zone; in 3 cases, access to the fracture site was performed to reposition the fragments; in 1 case, access to the fracture site was necessary to remove the migrated metal structure.

The average age of the patients was 48.6 years:

- in group I - 53.7;
- in the II group - 43.5 years.

Upon admission to the hospital, the primary method for stabilizing fresh fractures was skeletal traction - 25 patients or a rod unilateral apparatus for extrafocal fixation (EF) - 2 patients. Final stabilization of a fresh fracture was performed:

- during the first day from the moment of injury - 1 patient;
- for 2–5 days - 2 patients;
- 6-10 days - 20 patients;
- 11–20 days - 4 patients, which was due to the need to stabilize the general condition of the patient.

Fracture fixation surgical technique

Outer osteosynthesis of metadiaphyseal fractures was performed using the MIPO technique - closed reposition of fragments under the control of an electron-optical converter (EOP), bridge fixation of the fracture from a short (up to 5 cm) lateral approach.

In type 33 fractures, the fragments were openly repositioned through the lateral parapatellar approach.

In case of type 33-C fractures, revision of the knee joint, reposition of intra-articular fragments, fixation of them with spongy screws, after which a plate was installed and the intra-articular block was fixed to the diaphyseal fragment, was performed. The axial parameters were verified under the control of the eye.

In 13 cases the LCP-DF plate was used, in one case the LISS plate was used. The plates were fixed along the lateral surface of the femur.

Blocking intramedullary osteosynthesis (BIOS) of the distal femur in all cases was performed with a retrograde nail from the medial parapatellar approach without reaming the bone marrow canal.

For fractures 33-C, revision of the knee joint, condyles juxtaposition, and rod placement were performed.

For intra-articular fractures, a reconstructive rod with tie bolts was used to fix the condyles.

The insertion point of the rod is standard, in the intercondylar zone of the thigh. In one case, a primary dynamic BIOS was performed (fracture 32-A3), in all the others - a static BIOS of the distal femur with subsequent dynamization.

Research methods

In our work, we used X-ray and clinical research methods. Treatment results were assessed using the Mattis scale [3] and the Knee Score [5, 10]. The observation period for patients is up to 2 years after the operation.

In the scale for assessing the function of the knee joint after injury (Knee Score), the following indicators are considered:

- 1) subjective - pain at rest, prolonged forced position;
- 2) objective - range of motion, presence and severity of contracture, axis of the limb;
- 3) functional - the distance of movement, walking on stairs, the presence of external support when moving.

The orthopedic regimen included early active function in the knee joint (3-4 days after surgery), limited load on the limb.

Medical therapy is standard for trauma patients with fractures of the lower extremities.

Results and its discussion

Evaluation of treatment results according to the Matthis scale and the Knee Score scale showed similar treatment results in the first and second groups of patients.

Both tables show approximately similar treatment results in both groups. This is due to the fact that the analysis of the results in each of the groups was carried out without dividing by the type of fracture. The most interesting are indicators of treatment outcomes with a separate distribution by types of fractures.

With type 32 fractures, an equally good recovery of joint function is observed, while, due to the better biomechanics of the intraosseous fixator, in these types of fractures in group II, an earlier activation of patients was observed, which had a positive effect on the dynamics of restoration of the activity of the injured limb.

Fractures of type 33-A in patients treated with external osteosynthesis have a high percentage of recovery of the knee joint function - 79%; with the use of an intraosseous fixator, lower indicators of the function of the knee joint are noted - 62%.

The results of restoration of the function of the knee joint in case of comminuted intra-articular fractures of type 33 - C with the use of an extra-bone fixator are about 70%, and with the use of an intraosseous nail - only 41%.

Complications during treatment

In group I, X-ray fusion of a fracture against the background of osteosynthesis with a plate with angular stability was obtained in 12 (85.7%) patients out of 14 at standard times, which are common for a fracture of this type and localization.

The following complications were observed in 2 patients:

- one patient (7.1%) had a fatigue fracture of the plate, for which later reosteosynthesis was performed with the same fixator;
- in another patient (7.1%) with a concomitant diagnosis of type I diabetes mellitus, the locking screws were pulled out of the bone, which caused a change in the fixation technique.

Both cases occurred among patients of the older age group (71 and 72 years) with the same fractures - type 32-B1 at 3.5 and 6 months after osteosynthesis. There was no secondary displacement or disruption of the limb axis.

In 2 patients, it was difficult to remove the extraal fixator (the effect of “cold welding” between the heads of the locking screws and the holes of the plate).

In group II, X-ray analysis showed fracture union in all patients. In 12 (92.3%) cases, fusion occurred at the usual time for a fracture of this type and localization.

One patient (7.7%) with a double hip fracture had a rod fracture followed by reosteosynthesis.

In 4 patients, complications were observed that did not lead to an increase in the timing of the fracture union:

- in 1 (7.7%) patient, it was not possible to achieve anatomical articular reduction (fracture 33 - C2.3);

- 1 (7.7%) patient had an iatrogenic fracture of the proximal fragment (fracture 32 - A1.3);

- in 2 (15.4%) patients (fractures 33 - C), secondary displacement of the distal fragment with a violation of the femoral axis occurred.

Infectious complications were not observed in any group of patients.

When choosing a method for treating periarticular fractures, it should be borne in mind that the intrasosseous canal of the femur has an elliptical shape - the sagittal diameter is larger than the frontal one. With this shape of the canal, the intrasosseous rod is in close contact with the lateral and medial walls of the canal, which ensures stability of fixation. In the distal part of the femur, the intrasosseous canal is widened in the frontal plane, which prevents tight contact of the nail with the bone in this zone. Thus, with intramedullary osteosynthesis of fractures of the distal femur, it becomes necessary to increase the rigidity of fixation of the distal fragment. In our study, for this purpose, we used universal retrograde rods, in which the distal blocking was performed with ChM® tie bolts and Targon retrograde rods with the possibility of fixing the distal fragment with 4 screws in one plane from Magma-Sich®. There are other designs of intramedullary fixators that allow to achieve an increase in rigidity in the distal femur, but these fixators are not officially presented in Uzbekistan.

When comparing the results of surgical treatment with the use of various fixators, it was noted that with extra-articular metadiaphyseal fractures of type 32, fixation of the BIOS is more stable and allows earlier activation of patients. The use of a plate with angular stability in this case is less preferable, since the biomechanical axis of the limb passes medial to the plate, this somewhat limits the early loading, and if an attempt is made to load early before consolidation, fractures of the fixators are possible.

Intra-articular fractures of type 33-C require anatomical comparison of the intra-articular components of the fracture and rigid fixation of the articular block to the diaphyseal fragment. Analysis of complications in group II (in 2 patients there was a secondary displacement of the distal fragment with a violation of the femoral axis; in one patient the congruence of intra-articular fragments was not initially achieved; in all cases there were 33-C intra-articular fractures) suggests that in the case of fractures 33-C it is more expedient to perform an open reduction, achieve anatomical comparison of the fragments under the control of the eye, and fix the fracture with a plate. This provides greater stability of the osteosynthesis to secondary displacement of fragments, despite the possibility of some rod designs using tie bolts to fix the femoral condyles.

The most ambiguous is the algorithm for fixing type 33-A fractures. Currently, there is no consensus in the domestic and foreign literature regarding the fixation of fractures of the distal femur: traumatologists were conditionally divided into supporters of osteosynthesis with plates [8, 11, 14] and BIOS with a retrograde rod [7, 9, 12, 15].

In our study, we observed 4 patients with 33-A fractures in group I and 6 patients in group II. Fracture union was obtained in all cases.

There were no complications during plate osteosynthesis. Postoperative follow-up for up to 2 years showed restoration of function in the knee joint in all patients.

In the group where the osteosynthesis was performed with an intramedullary nail, in 1 case there was a secondary displacement of the fragments, the femur axis was broken. In our opinion, given the absence of special rods with increased rigidity of fixation of the distal fragment, it is more expedient to fix these fractures with plates with angular stability using minimally invasive technique (MIPO).

Conclusions

1. In extra-articular metadiaphyseal fractures of type 32, fixation of the BIOS with indirect fracture reduction is more preferable due to greater stability and less trauma to soft tissues.

2. In case of type 33-C intra-articular fractures, it is necessary to achieve anatomical reduction of articular fragments, preservation or restoration of the axial relationship of the femur and tibia, which is better achieved with open reduction. Fixation of intra-articular fractures should be carried out with an extra-bone fixator. In the treatment of elderly patients and in the case of unstable fractures (33 - C2, C3), it is more expedient to use a plate with angular stability to prevent secondary displacement of fragments.

3. Fixation of fragments in type 33 - A fractures with an angularly stable extrasteel plate (especially when using the MIPO technique) provides greater stability and fewer complications.

References

1. V.R. Akramov, Sh.Sh. Akhmedov, B.U. Khamraev, A.A. Teshaev E.M. Khayatov, U. U. Radjabov., Endoprosthetics of the hip joint for fractures of the femoral neck // Problems of biology and medicine.- Samarkand No. 3 (96) 2017. pp. 23-26. [in Russian]
2. Akramov V.R. Features of hip arthroplasty in case of anatomical disorders of the acetabulum // "Bulletin of the association of doctors of Uzbekistan" Uzbekistan, Tashkent № 3 - 2011, Pages 94-97. [in Russian]
3. Akhmedov Sh.Sh., Khamraev A.Sh., Akramov V.R., Tugizov B.E., Khamraev B.U. Prophylactic features of pulmonary artery thromboembolism after hip arthroplasty in dysplastic coxarthrosis // "Bulletin of the association of doctors of Uzbekistan" Uzbekistan Tashkent № 1 (98) - 2020, pp. 42-47, pp. 42-47. [in Uzbek]
4. Akramov V.R. Some problems of hip joint replacement previously operated on. // "Bulletin of the association of doctors of Uzbekistan" Uzbekistan Tashkent № 2 - 2011, pp. 110-113. [in Russian]

5. Akramov V.R., Akhmedov Sh.SH., Khamraev B.U.- (Hip replacement in femoral neck fractures) // "Problems of biology and medicine" Uzbekistan, Samarkand No. 3 - 2017 (96), pp.23-26 [in Russian]
6. Akramov V.R., Sh.Sh., Khamraev A.Sh., Khamraev B.U. – (Total hip replacement and prevention of possible complications) // "A new day in medicine" Uzbekistan. Tashkent, No.4 (20) 2017, pp.56-58. [in Russian]
7. Akramov V.R., Akhmedov Sh.SH., Khamraev A.SH., Khamraev B.U. - (Hip replacement in degenerative-dystrophic diseases in adults) // "Bulletin of the association of doctors of Uzbekistan" Uzbekistan, Tashkent No. 2 - 2018, pp.42-44. [in Russian]
8. SH.SH.Akhmedov, A.SH.Khamraev, V.R.Akramov, B.U.Khamraev, A.A.Teshaev, A.U. Gaffarov The arthroplasty of the hip at fracture of a neck of a femur // "A new day in medicine "Uzbekistan.Tashkent, No. 1 (25) 2019, pp. 5-7.
9. V.R.Akramov, B. A.SH.Khamraev, SH.SH.Akhmedov, B.U.Khamraev The Arthroplasty Of The Hip At Fracture Of A Neck Of A Femur // European Journal of Business & Social Sciences., ISSN: 2235-767X., Volume 07 Issue 05., May 2019.
10. V.R.Akramov, B.A.SH.Khamraev, SH.SH.Akhmedov, B.U.Khamraev // Prevention Of Possible Complications Before And After Total Endoprotésization Of The Combin) European Journal of Business & Social Sciences., ISSN: 2235-767X., Volume 07 Issue 05., May 2019.
11. Akhmedov Sh.Sh., Khamraev A.Sh., Vazina G.P., Akramov V.R., Khamraev B.U., Tugizov B.E., THE Peculiarities of prophylaxis of pulmonary thromboembolism after total hip endoprosthesis in dysplastic coxarthrosis // "New day in medicine" Uzbekistan.Tashkent, №2 (30) 2020, pp. 53-56.
12. Nurullaev S.O; Akhmedov Sh.Sh; Akramov V.R; Khamraev A.Sh; Khamraev B.U; Mirzamurodov Kh.Kh Our experience in the treatment of grade I-II gonarthrosis with hyaluronic acid preparations // Academicia An International Multidisciplinary Research Journal, Vol. 10 Issue 12, December 2020.

13. Asilova Saodat Ubayevna., Akramov Voxid Rustamovich., Akhmedov Shamshod Shavkatovich., Mirzamurodov Khabibjon Khalimovich (Tashkent, Uzbekistan). // Polish science journal., Issue 12 (33) Part 2. Warsaw, Poland Wydawnictwo Naukowe "iScience" 2020. [in Russian]
14. Yuldashev R.M., Mardanov J.J. Vertebroplasty in the surgical treatment of spinal tumors - evaluation of treatment results. Scientific and practical journal // "Doctor-graduate student" No. 4 (53). 2012.S. 9-13. [in Russian]
15. Yuldashev R.M., Mardanov Zh.Zh. Our experience in the treatment of patients with extradural tumors of the spinal cord. // Journal of Theoretical and Clinical Medicine. 2013. - No. 3.- P.100-102. [in Russian]
16. Kariev G.M., Mardanov J.J., Norov A.U. Pathogenesis of metastasis in the spine. // Scientific and Practical Journal of Neurology 2014.-№4.- P. 37-39. [in Russian]
17. Mardanov J.J., Zikriyayev N.N., Rakhmatov K.R., Rajabov M.M. Quality of life in patients with extradural tumors of the spinal cord. // Journal of Surgery of Uzbekistan 2016.-№4-C-25-27. [in Russian]
18. Mardanov J.J. Surgical treatment of spinal hemangiomas. // Russian neurosurgical journal named after professor A.L. Polenov. 2014 volume-№4. –S.- 343-345. [in Russian]
19. Mardanov J.J.The result of surgical treatment of pathological pathological fracture during extradural tumor of spinal cord. // European Sciences Review 2014.- №3-4 - S.-21-24.
20. Mardanov J.J. Back - side access during surgical treatment of extradural tumors of the spinal cord. // Questions of science and education. 2021. - № 22 C. - 147. [in Russian]
21. Mardanov J.J. Surgical treatment of extradural tumors of the spinal cord. // Questions of science and education. 2021. - № 26 S. - 157. [in Russian]

22. Khamraev B.U., Akhmedov Sh.Sh. Two-stage revision hip replacement patients with severe acetabulum defect (case report) // Asian journal of Pharmaceutical and biological research. Volume 10 Issue 2 MAY-AUG 2021 P. 35-41.
23. Khamraev B.U., Akhmedov Sh.Sh., Our experience of treatment of femur fractures by the method of intramedullar locking osteosynthesis.// Asian journal of Pharmaceutical and biological research. Volume 10 Issue 2 may-aug 2021. P. 42-46.
24. BU Khamraev, BP Akramov. Program for expressing the method of treatment by the method of blocking intramedullary osteosynthesis for a fracture of the femur // Certificate of official registration of a computer program. Agency for Intellectual Property of the Republic of Uzbekistan. 2019. [in Russian]
25. Nematov Dilshod Amrilloevich. Features of the application of external osteosynthesis in gonarthrosis // Asian journal of Pharmaceutical and biological research. Volume 10 Issue 2 MAY-AUG 2021. P. 26-30.
26. Ziyadullaev Abdusalom Khabibulla oglu. Evaluation of the future results of application of arthro-medullary bypassing in gonarthrosis // Asian journal of Pharmaceutical and biological research. Volume 10 Issue 2 MAY-AUG 2021. P. 31-34.
27. Mirzamurodov Habibjon Halimovich, Nurulloev Sukhrob Ozodovich. Improvement of surgical treatment of patients with combined degenerative-dystrophic pathology of the hip joint and spine with prevalence of manifestations of coxarthrosis // British Medical Journal Volume-1, No 2., 2021. P.180-187.
28. Nurulloev S.O., Mirzamuradov H.H. Morphological Changes In Bone Tissue In Chronic Osteomyelitis On The Background Of Application Of Plate Concentrate // The American Journal of Medical Sciences and Pharmaceutical Research Volume 3 Issue 04, 2021. P. 160-164.
29. Mirzamurodov H.H. New approaches to treatment of patients with coxovertebral syndrome // Asian journal of Pharmaceutical and biological research. Volume 10 Issue 2 MAY-AUG 2021. P. 9-19

30. Mirzamurodov Kh.Kh., Akhmedov Sh.Sh., Nuruloev S.O., Ziyadullaev A.Kh., Nematov D.A. Optimization of total hip arthroplasty in dysplastic coxarthrosis // New day in medicine. 4 (32) 2020 P. 667-672. [in Russian]
31. Mirzamurodov H. Kh., Khodzhanov I. Yu., Nurulloev S.O. Complex conservative therapy for hip-spine syndrome // International Scientific and Educational electronic journal "Education and science in the xxi century" 2021. Issue No. 12 (volume 2). P. 1438-1439. [in Russian]
32. Nurulloev S.O., Mirzamuradov Kh.Kh. Our experience in the treatment of degree I-II gonarthrosis with drugs hyaluronic acid // Innovation in the modern education system. 2021. Part 5, Issue 1. P. 546-548. [in Russian]
33. Sulaymanova Gulnoza Tulkindzanovna, Amonov Muhammad Komilovich. Regional Causes Of Iron Deficiency Anemia, Pathogenesis And Use Of Antianemic Drugs. // The American Journal of Medical Sciences and Pharmaceutical Research (ISSN – 2689-1026) Published: April 30, 2021. P. 165-170.
34. Sulaymonova Gulnoza Tulkinjanovna, Raufov Alisher Anvarovich. The influence of deficiency of microelements in children with bronchial hyperreactivity // ACADEMICIA: An International Multidisciplinary Research Journal (ISSN: 2249-7137) Published: Vol. 10, Issue 4, April 2020 | Pages: 846-853
35. Sulaimonova G.T., Amonov M.K., Rakhmonova K.E. The frequency of detection of risk factors for chronic kidney disease in the rural population. // Bulletin of Science and Education № 24 (102). Part 2. 2020. pp. 79-85. [in Russian]
36. Orziev Zavkidin Mansurovich, Suleimanova Gulnoza Tulkinzhanovna. Regional causes of iron deficiency anemia in women of fertile age. // Electronic scientific journal "Biology and Integrative Medicine" №4 – April (21) 2018. Pp. 74-82. [in Russian]
37. ZM Orziev, GT Sulaimonova. Analysis of modern ideas on the formation of critical periods in the event of iron deficiency anemias (review) // Bulletin of the

Council of Young Scientists and Specialists of the Chelyabinsk Region Volume 5, No. 4, 2017. P. 17-25 [in Russian]

38. Abdullaeva U.K. Predicting the risk of atrophic transformation in chronic gastritis using serum pepsinogen // World journal of pharmaceutical research, Faculty of Pharmacy Medical University, Bulgaria, Vol. 8, Iss. 13, 2019, P. 219-228.

39. Abdullaeva U.K., Sobirova G.N., Karimov M.M., Aslonova I.J. The prevalence and possibilities of prevention of noncardial gastric cancer in the Bukhara region // American journal of medicine and medical sciences, 2020, 10(9), P. 679-681.

40. Sobirova G.N., Abdullaeva U.K., Nosirova M.S., Aslonova I.J. Evaluation of the gastrointestinal mucosa by the OLGA system in chronic atrophic gastritis // Journal of critical reviews, Kuala Lumpur, Malaysia, Vol. 7, Iss. 2, 2020, P. 409-413.

41. Karimov M.M., Sobirova G.N., Abdullaeva U.K., Aslonova I.Zh., Tulyaganova F.M. Possibilities of serological diagnosis of atrophic processes of the gastric mucosa // European Journal of Molecular & Clinical Medicine Vol. 7, Iss. 11, 2020, P. 2955-2960.

42. Karimov M.M., Sobirova G.N., Abdullaeva U.K. Chronic gastritis and carcinogenesis issues // Herald of Pancreatic Club, 2019. Iss. 45 (4). P. 65-70. [in Russian]

44. Sobirova G.N., Abdullaeva U.K. Immunopatogenesis of chronic gastritis and its role in carcinogenesis // Journal of Biomedicine and Practice, 1 (4). P. 40-44.

45. Karimov M.M., Sobirova G.N., Abdullaeva U.K., Aslonova I.Zh., Tulyaganova F.M. Possibilities of Serological Diagnosis of Atrophic Processes of the Gastric Mucosa // Annals of the Romanian Society for Cell Biology, , Vol. 25, Issue 1, 2021, Pages. 6168 – 6174.

46. Abdullaeva U.K., Shadjanova N.S. Using the OLGA system in chronic atrophic gastritis // New day in medicine, 2020, №2, P. 9-12.

47. Abdullaeva U.K. The value of interactive teaching methods in improving the level of clinical knowledge of students // Medical education and professional development. 2019, №1 (33), P. 29-32. [in Russian]
48. GN Sobirova, UK Abdullaeva Chronic gastritis and carcinogenesis issues // Central Asian Problems of Modern Science and Education. 2019, Iss. 4, №2, P. 159-172
49. Orziev Zavkiddin Mansurovich, Abdullaeva Umida Kurbanovna, Nurkhanova Nilufar Odiljonovna Study of the effectiveness of cholelitholytic therapy in patients with cholelithiasis, taking into account the type of violation of the contractility of the gallbladder // Science of the Young - Eruditio Juvenium. 2015. №4. P. 50. [in Russian]
50. Karimov M.M., Sobirova G.N., Abdullaeva U.K., Aslonova I.Zh., Tulyaganova F.M. Serological Diagnosis of Atrophic Processes of the Gastric Mucosa // The American Journal of Medical Sciences and Pharmaceutical Research, Vol. 2, Issue 12, 2020, Pages. 118-124
51. DB Mirzaeva, UK Abdullaeva, RR Boboeva The importance of interactive teaching methods in improving the level of clinical knowledge of students // Central Asian Problems of Modern Science and Education Vol. 4, Issue 2, 2019, Pages. 159-166
52. MM Karimov, ST Rustamova, ZhA Ismailova, UK Abdullaeva, ZZ Saatov Diagnostic efficacy of C14 breath test in Helicobacter pyloriosis // Cardiovascular therapy and prevention. 2019. Vol. 18, Issue S1, P. 85-86
53. Abdullaeva U.K., Jalolova V.Z. Study of the effectiveness of cholelitholytic therapy in patients with cholelithiasis, taking into account the type of violation of the contractility of the gallbladder // Bulletin of the Council of Young Scientists and Specialists of the Chelyabinsk Region. 2016. Vol. 5, Issue 4(15), P. 85-86 [in Russian]

54. Orziev Z.M., Abdullaeva U.K. Regional causes of extrahepatic "Subtransaminasemia" // Biology and integrative medicine 2016, №3. P. 28-40. [in Russian]
55. Abdullaeva U.K., Mirzaeva D.B. Regional prospects for metabolic therapy for stable senocardia Summary. // Bulletin of the South Kazakhstan Medical Academy 2019. P. 74-76 [in Russian]
56. Orziev Z.M., Abdullaeva U.K., Yuldasheva D.H. Method for early prediction of the efficiency of cholelytic therapy based on dynamic control of bild pH indicators in patients with cholelystone disease // Innovative development of modern science. 2014. P.76-79 [in Russian]
57. Orziev Z.M., Abdullaeva U.K. Relationship between the effectiveness of cholelitholytic therapy and the state of contractility of the gallbladder // Bulletin of the Council of Young Scientists and Specialists of the Chelyabinsk Region. 2015. №3(10) [in Russian]
58. Orziev Z.M., Abdullaeva U.K. The effectiveness of cholelitholytic therapy for cholelithiasis // Health is the basis of human potential: problems and ways to solve them. 2015. Iss. 10. №2. P. 610-612. [in Russian]
59. Hamroev Behzod Uktamovich, & Akhmedov Shamshod Shavkatovich. (2021). Optimization of blocking intramedullary osteosynthesis methods for femoral fractures // Asian Journal of Pharmaceutical and Biological Research 2231-2218, Volume 10 (Issue 3 Sept-Dec 2021), P. 29–43.